

FUNCTIONAL HEALTH LITERACY IN INDIVIDUALS WITH SPINAL CORD INJURY TREATED AT A REHABILITATION CENTER

Francine Aguilera Rodrigues da Silva¹; Cejane Oliveira Martins Prudente²;
Vanessa da Silva Carvalho Vila³; Katarinne Lima Moraes⁴;
Maria Alves Barbosa⁵; Celmo Celeno Porto⁶

Highlights: (1) Functional health literacy was adequate in 70.6% of individuals. (2) Age, income and having a partner: indicators of better health literacy. (3) Rehabilitation time and education influence the results of health literacy.

PRE-PROOF

(as accepted)

This is a preliminary, unedited version of a manuscript accepted for publication in Revista Contexto & Saúde. As a service to our readers, we are making this initial version of the manuscript available as accepted. The article will still be reviewed, formatted, and approved by the authors before being published in its final form.

<http://dx.doi.org/10.21527/2176-7114.2025.50.15306>

How to cite:

da Silva FAR, Prudente COM, Vila V da SC, Moraes KL, Barbosa MA, Porto CC. Functional health literacy in individuals with spinal cord injury treated at a rehabilitation center. Rev. Contexto & Saúde, 2025;25(50): e15306

¹ Universidade Federal de Goiás – UFG. Goiânia/GO, Brasil. Centro Universitário Alfredo Nasser - UNIFAN. Aparecida de Goiânia/GO, Brazil. <https://orcid.org/0000-0003-1368-7924>

² Pontifícia Universidade Católica de Goiás - PUC Goiás. Goiânia/GO, Brasil. Universidade Estadual de Goiás – UEG. Goiânia/GO, Brazil. <https://orcid.org/0000-0001-6499-3011>

³ Pontifícia Universidade Católica de Goiás - PUC Goiás. Goiânia/GO, Brazil. <https://orcid.org/0000-0002-1785-8682>

⁴ Universidade de Brasília – UnB. Brasília/DF, Brazil. <https://orcid.org/0000-0001-6169-0461>

⁵ Universidade Federal de Goiás – UFG. Goiânia/GO, Brazil. <https://orcid.org/0000-0002-0861-9655>

⁶ Universidade Federal de Goiás – UFG. Goiânia/GO, Brazil. <https://orcid.org/0000-0001-5860-9490>

FUNCTIONAL HEALTH LITERACY IN INDIVIDUALS WITH SPINAL CORD INJURY TREATED AT A REHABILITATION CENTER

ABSTRACT

This study aims to analyze the functional health literacy of individuals with spinal cord injury treated at a rehabilitation center. We conducted this analytical cross-sectional study at a rehabilitation and readjustment center, including 85 individuals diagnosed with spinal cord injury. We collected data on sociodemographic, clinical, and functional profiles. The Short Test of Functional Health Literacy in Adults (S-TOFHLA) was used to assess functional health literacy. Most participants were male, aged between 18 and 39 years, with paraplegia caused by traumatic etiology. Analysis of functional health literacy showed that 70.6% of individuals with spinal cord injury had adequate literacy levels. Better functional health literacy outcomes were associated with younger age, higher income, longer duration of rehabilitation, higher educational levels, and having a partner. In conclusion, the services provided and the organizational structure of reference centers rehabilitating individuals with spinal cord injury may contribute to improved functional health literacy. Identifying related variables and analyzing the functional dimension of health literacy by multidisciplinary teams during rehabilitation are essential tools for enhancing the health outcomes of individuals with spinal cord injury.

Keywords: health literacy, spinal cord, patient care team, health education.

INTRODUCTION

Spinal cord injury (SCI) is characterized by damage to the spinal cord, which can cause motor, sensory, and autonomic impairments¹. SCI results in various sequelae that significantly disrupt affected individuals' routines and quality of life, necessitating continuous learning and adaptation².

According to the 2019 Global Burden of Disease Study, there were an estimated 0.9 million incident cases and 20.6 million prevalent cases of SCI worldwide. The incidence has increased over the past 30 years³. Projections for 2030 indicate a rising trend in age-standardized incidence rates for both sexes⁴.

Individuals affected by SCI require specific guidance, making it essential for them to participate in the rehabilitation process. Adherence to this process depends on collaboration between patients with SCI, their families, and healthcare professionals. Clear communication, appropriate treatment, and early initiation of rehabilitation are fundamental to restoring

FUNCTIONAL HEALTH LITERACY IN INDIVIDUALS WITH SPINAL CORD INJURY TREATED AT A REHABILITATION CENTER

autonomy and decision-making control to individuals with SCI⁵.

Health literacy (HL) plays a critical role in developing self-management skills, making therapeutic decisions, reducing complications, and decreasing treatment costs⁶. Initially, HL was associated with reading and understanding health materials; however, more recent definitions encompass multidimensional skills related to health promotion, influenced by social and demographic factors⁷.

The level of HL can be categorized into three dimensions: functional health literacy (FHL), interactive health literacy, and critical health literacy. The first dimension involves the ability to read basic health information; the second relates to the capacity to seek and discuss more complex information; and the third pertains to autonomy in making informed health decisions⁸.

The rehabilitation of individuals with SCI is a collaborative process requiring constant information exchange and the development of trust between professionals and patients⁹. Inadequate FHL can directly affect the rehabilitation process, emphasizing the need for health information to be accessible, comprehensible, and applicable to everyone, particularly to at-risk groups such as individuals with disabilities^{10,11}.

This study focused on analyzing FHL, which relates to the ability to read, comprehend texts, and understand numerical information in the healthcare context, enabling its application in daily life¹⁰. FHL encompasses the skills needed to seek, understand, and use health information to make decisions and adopt healthy behaviors, such as adhering to treatments, participating in prevention programs, and changing habits⁸.

Given that FHL may impact the autonomy, independence, and readaptation of individuals with SCI to their new life circumstances¹¹, its assessment becomes particularly relevant. In this context, the present study aimed to analyze the functional health literacy of individuals with spinal cord injury treated at a rehabilitation center.

METHODS

We conducted this cross-sectional analytical study at a rehabilitation and adaptation center in Goiânia, the capital of Goiás state, Brazil. The Ministry of Health recognizes the center as a Specialized Rehabilitation Center (CER IV). It exclusively operates under the Unified Health System (SUS)—providing rehabilitation services for individuals with

FUNCTIONAL HEALTH LITERACY IN INDIVIDUALS WITH SPINAL CORD INJURY TREATED AT A REHABILITATION CENTER

physical, auditory, visual, and intellectual disabilities—and serves as a reference center for SCI care.

The inclusion criteria were: patients undergoing treatment at the rehabilitation center, with a diagnosis of traumatic or non-traumatic SCI documented in their medical records, aged 18 years or older, of any sex, with any level or duration of SCI, with at least one year of formal education or self-reported informal acquisition of reading skills (individuals who did not attend school but could read were considered to have one year of education), and without other associated neurological and/or orthopedic injuries.

Individuals were excluded if they had documented or reported neurological or cognitive impairments that could interfere with understanding the study instruments, severe visual or hearing impairments that could not be mitigated with assistive technologies, and incomplete medical records.

At the time of data collection, 92 patients with SCI were receiving care at the rehabilitation center. Based on the inclusion and exclusion criteria, 87 patients remained eligible, and after contact, two declined participation. Thus, 85 patients were interviewed between October 2021 and January 2022.

The research team, trained in data collection procedures, utilized the following tools: the Short Test of Functional Health Literacy in Adults (S-TOFHLA), electronic medical records for completing the clinical and functional profile form, and a questionnaire for sociodemographic characterization.

The S-TOFHLA is a shortened version of the Test of Functional Health Literacy in Adults (TOFHLA), validated and adapted to the Brazilian context¹². This instrument consists of 36 items evaluating reading comprehension and four items assessing numeracy skills related to health topics, such as diagnostic test instructions, consent forms, appointment scheduling, and prescription labels^{12,13}. The maximum time allowed for completing the tasks is 12 minutes, after which participants are asked to stop¹⁴. Participants are not informed of this time limit beforehand. Reading comprehension scores range from 0 to 72 points, with two points assigned for each correctly completed blank space. Numeracy items total 28 points, with seven points awarded for each correct response. The test's total score is 100 points¹³.

Based on performance, participants are classified into three levels of FHL: inadequate (0–53 points), marginal or borderline (54–66 points), and adequate (67–100 points)¹⁵.

Medical record data included injury duration, rehabilitation duration, etiology, injury

FUNCTIONAL HEALTH LITERACY IN INDIVIDUALS WITH SPINAL CORD INJURY TREATED AT A REHABILITATION CENTER

level, type of impairment, and SCI classification based on the American Spinal Injury Association (ASIA) Impairment Scale (AIS), categorized as A, B, C, D, and E. AIS A represents complete SCI, where no sensory or motor function is preserved below the neurological level. AIS B indicates incomplete SCI, with no motor activity below the neurological level but some preserved sensory function. AIS C and D signify reduced motor activity below the neurological level, while AIS E indicates normal sensory and motor function¹⁶.

The sociodemographic profile questionnaire, developed by the researcher, was based on prior studies on FHL in chronic diseases and included data on sex, age group, marital status, place of origin, family income, and education level¹⁷⁻¹⁹.

Patient interviews were conducted in the waiting room of the institution's adult neurofunctional physical therapy outpatient clinic, lasting 40–50 minutes. During the S-TOFHLA, reading comprehension was allocated seven minutes, while numeracy tasks had a maximum of five minutes²⁰.

We analyzed data using the Statistical Package for the Social Sciences (SPSS) version 26.0. Data normality was assessed using the Shapiro-Wilk test. Sociodemographic, clinical, and FHL profiles were described using absolute frequency (n), relative frequency (%), means, standard deviations, medians, and ranges for continuous variables.

Exploratory analysis of FHL and patient profiles utilized Spearman's correlation, Mann-Whitney tests, and Kruskal-Wallis tests, followed by Nemenyi post hoc tests.

We included variables with a p-value < 0.20 in the linear regression from the initial exploratory analysis to identify potential predictors. Finally, multiple linear regression analysis was performed between the total S-TOFHLA score and predictor variables using the Stepwise method. The significance level was set at 5% ($p < 0.05$).

The study adhered to the recommendations of Resolution 466/12 of the National Health Council and was approved by the Research Ethics Committee (approval number: 5.015.904).

**FUNCTIONAL HEALTH LITERACY IN INDIVIDUALS WITH SPINAL
CORD INJURY TREATED AT A REHABILITATION CENTER**

RESULTS

Eighty-five individuals diagnosed with SCI were evaluated, including 22 females (25.9%) and 63 males (74.1%). The majority were aged between 18 and 39 years and did not have a partner (51.8% never married and 9.4% widowed or divorced). The sociodemographic variables evaluated are described in Table 1.

Table 1. Sociodemographic profile of 85 patients with spinal cord injury treated at a rehabilitation center, Goiânia, Goiás, 2022

	N	%
Sex		
Female	22	25.9
Male	63	74.1
Age group		
18 to 39	46	54.1
40 to 59	31	36.5
60 to 71	8	9.4
Having a partner		
Yes	33	38.8
No	52	61.2
Place of origin (Goiás)		
Metropolitan area (Goiânia)	60	70.6
Non-metropolitan area	25	29.4
Family income		
Up to 2 minimum wages	59	69.4
2 to 4 minimum wages	21	24.7
4 to 10 minimum wages	5	5.9
Education level		
1–9 years of schooling	38	44.7
10–12 years of schooling	40	47.1
> 12 years of schooling	7	8.2

n = absolute frequency; % = relative frequency

**FUNCTIONAL HEALTH LITERACY IN INDIVIDUALS WITH SPINAL
CORD INJURY TREATED AT A REHABILITATION CENTER**

Regarding the injury classification, a predominance of complete injuries, categorized as AIS A (42.4%), was observed. Most cases had a traumatic origin, with an average injury duration of 55.5 months and a rehabilitation duration of 38.06 months. The coefficient of variation (CV) for injury duration was 110.51%, and for rehabilitation duration, it was 97.23%, indicating considerable heterogeneity among patients in these aspects. The clinical profile of the patients is presented in Table 2.

Table 2. Clinical profile of 85 patients with spinal cord injury treated at a rehabilitation center, Goiânia, Goiás, 2022

	Mean \pm DP	Median (Minimum - Maximum)
Injury duration (months)	55.55 \pm 61.37	33.0 (1.0 - 330.0)
Rehabilitation duration (months)	38.06 \pm 37.01	25.0 (1.0 - 181.0)
	N	%
Injury etiology		
Motor vehicle accidents	38	44.7
Shallow water diving	4	4.7
Gunshot wounds	15	17.6
Falls from heights	9	10.6
Other (non-traumatic)	19	22.4
Injury classification (AIS)		
A	36	42.4
B	7	8.2
C	23	27.1
D	19	22.4
Level of injury		
Cervical	29	34.1
Thoracic	46	54.1
Lumbar	10	11.8
Sacral	-	-
Type of impairment		
Tetraplegia	29	34.1
Paraplegia	56	65.8

n = absolute frequency; % = relative frequency; SD = standard deviation

**FUNCTIONAL HEALTH LITERACY IN INDIVIDUALS WITH SPINAL
CORD INJURY TREATED AT A REHABILITATION CENTER**

The majority of respondents, 60 (70.6%), demonstrated adequate functional health literacy (FHL), while 13 (15.3%) had borderline FHL, and 12 (14.1%) exhibited inadequate FHL. The mean score for the total S-TOFHLA was 79.56 ± 21.34 .

Regarding reading comprehension, the mean score was 58.40 ± 14.55 (range: 20.00–72.00), whereas for numeracy, it was 21.18 ± 9.44 (range: 0.00–28.00).

Among participants with inadequate FHL, nine were unable to complete the entire instrument. These participants failed to fill in all the blanks for reading comprehension, exceeding the seven-minute time limit, at which point the test was stopped.

Table 3 compares the total FHL score with the profiles of patients with SCI. A significant difference ($p < 0.05$) was observed for the variables age group, having a partner, income, and education level. Younger individuals, those with a partner, higher income, and higher education levels achieved better total S-TOFHLA scores.

Table 3. Comparison of functional health literacy with the profile of 85 patients with spinal cord injury treated at a rehabilitation center, Goiânia, Goiás, 2022

	Total S-TOFHLA score
Age group**	p = 0.03
18–39	$85.91 \pm 16.36^\dagger$
40–59	74.61 ± 21.68
60–71	62.25 ± 31.88
Sex*	p = 0.25
Female	74.36 ± 24.22
Male	81.38 ± 20.14
Having a partner*	p = 0.04
Yes	84.91 ± 16.98
No	76.17 ± 23.22
Place of origin (Goiás)*	p = 0.34
Metropolitan area (Goiânia)	80.70 ± 21.51
Non-metropolitan area	76.84 ± 21.13
Family income**	p = 0.01
Up to 2 minimum wages	74.63 ± 23.13
2 to 4 minimum wages	$89.57 \pm 10.87^\dagger$

**FUNCTIONAL HEALTH LITERACY IN INDIVIDUALS WITH SPINAL
CORD INJURY TREATED AT A REHABILITATION CENTER**

4 to 10 minimum wages	95.80 ± 1.79†
Education level**	p = 0.02
1–9 years of schooling	73.18 ± 24.96
10–12 years of schooling	83.60 ± 16.54†
> 12 years of schooling	91.14 ± 15.32†
Injury etiology**	p = 0.22
Motor vehicle accidents	81.37 ± 22.04
Shallow water diving	93.00 ± 10.00
Gunshot wounds	80.00 ± 15.38
Falls from heights	76.78 ± 23.64
Other (non-traumatic)	74.11 ± 24.42
Injury classification (AIS)**	p = 0.48
A	80.08 ± 22.09
B	76.57 ± 14.70
C	82.74 ± 21.33
D	75.84 ± 22.76
Type of impairment*	p = 0.42
Tetraplegia	76.93 ± 25.67
Paraplegia	80.93 ± 18.84

* Mann-Whitney; ** Kruskal-Wallis; † Nemenyi (Mean ± Standard deviation)

The Spearman correlation results indicated a relationship between injury duration, rehabilitation duration, and FHL. Injury duration was positively correlated with the reading comprehension component ($p = 0.02$) and rehabilitation duration ($p = 0.03$). Moreover, injury duration also showed a positive correlation with the total S-TOFHLA score ($p = 0.04$).

Based on the initial exploratory analysis, variables with a p -value < 0.20 (injury duration, rehabilitation duration, age group, having a partner, income, and education level) were selected for multiple linear regression analysis.

Table 4 presents the multiple linear regression analysis results between the total S-TOFHLA score and the predictor variables. Together, these variables explained 41% (r^2) of the total FHL score.

In the multiple linear regression model, age was negatively associated with FHL, indicating that younger individuals demonstrated higher FHL levels. On the other hand,

**FUNCTIONAL HEALTH LITERACY IN INDIVIDUALS WITH SPINAL
CORD INJURY TREATED AT A REHABILITATION CENTER**

higher income, longer rehabilitation duration, and higher education levels were positively associated with FHL. Additionally, having a partner significantly contributed to higher FHL levels in this sample (Table 4 and Figure 1).

Table 4. Results of the multiple linear regression analysis (Stepwise Method) between the total S-TOFHLA score and predictor variables for 85 patients with spinal cord injury treated at a rehabilitation center, Goiânia, Goiás, 2022.

	r^2	Beta	Standard error	<i>T</i>	<i>p</i>
Income		0.28	3.26	3.04	0.00
Age		-0.43	0.14	-4.82	0.00
Rehabilitation duration (months)	0.41	0.22	0.05	2.57	0.01
Education level		0.20	3.04	2.25	0.03
With a partner		0.20	3.96	2.19	0.03




Figure 1. Scatterplots of the total S-TOFHLA score and the predictor variables age and rehabilitation duration for patients with spinal cord injury treated at a rehabilitation center, Goiânia, Goiás, 2022.

FUNCTIONAL HEALTH LITERACY IN INDIVIDUALS WITH SPINAL CORD INJURY TREATED AT A REHABILITATION CENTER

DISCUSSION

FHL among individuals diagnosed with SCI attending the outpatient clinic of a rehabilitation reference center was adequate in 70.6% of participants. Since these patients were undergoing rehabilitation, we opted to evaluate FHL using the S-TOFHLA. This instrument measures an individual's ability to utilize reading, writing, and numeracy skills necessary for completing specific tasks or acquiring new knowledge, aspects that are essential in the rehabilitation process for individuals with SCI¹².

Functional independence following SCI is one of the primary goals of patient care. Identifying FHL levels is particularly important for adapting treatment strategies provided by healthcare professionals and empowering patients to participate in the planning and implementation of therapeutic processes involved in their care²¹.

Most participants were male, young, and never married, which aligns with epidemiological data on this condition^{22,23}. Studies indicate that men exhibit significantly higher SCI rates, which can be explained by more frequent risk behaviors, such as exposure to hazardous work environments and engagement in high-risk activities³.

Predictors of higher FHL levels included younger age, higher income, greater educational attainment, having a partner, and longer rehabilitation duration. In the literature, other factors have also been associated with FHL, including sex, skin color, life satisfaction, and self-rated health status^{6,24}.

Age was negatively associated with FHL, indicating that younger individuals exhibited better FHL. Conversely, higher income, greater educational attainment, and longer rehabilitation duration were positively associated with FHL.

Multiple linear regression analysis between the total S-TOFHLA score and predictor variables revealed that age was the strongest determinant of FHL levels in this population, consistent with findings from studies on other chronic conditions²⁵⁻²⁷.

According to the literature, among vulnerable groups—those more likely to exhibit low levels of FHL—older individuals stand out as a key demographic^{28,29}. Older adults may process information at a slower pace, and their reduced cognitive abilities can limit their capacity to engage with and understand health-related information³⁰. Additionally, older patients tend to struggle more with tasks requiring reasoning or deductions about presented information, suggesting a higher prevalence of low FHL in advanced age groups³¹.

FUNCTIONAL HEALTH LITERACY IN INDIVIDUALS WITH SPINAL CORD INJURY TREATED AT A REHABILITATION CENTER

Regarding income, higher income levels were associated with better total S-TOFHLA scores. A cross-sectional study conducted in China with 1,000 participants also found that higher income correlated with improved FHL³².

The associations observed between FHL and variables such as age, socioeconomic status, and skin color reinforce the principle that FHL is shaped by complex social structures beyond educational attainment alone³³.

Individuals with lower income often face greater challenges in accessing healthcare services and receiving health information, which results in poorer FHL levels and clinical outcomes²⁷.

In a study involving 107 outpatient SCI patients, 86% demonstrated adequate HL¹⁵. The authors noted that the observed FHL levels exceeded expectations, likely because the study was conducted in a private clinic with economically disadvantaged minorities but with higher educational levels.

Greater educational attainment among SCI patients was associated with better FHL scores. While the relationship between education and FHL has been demonstrated in other chronic diseases^{6,34}, it is important to note that higher educational levels do not guarantee adequate FHL. Even individuals with advanced education may struggle with medical terminology and procedures³⁶.

Patients and their families must comprehend the guidance healthcare professionals provide to make decisions that promote effective self-management of their condition. A qualitative study involving patients and caregivers in a rehabilitation setting identified that spouses often led efforts to obtain information about rehabilitation¹⁸.

Patients assisted by their partners may find it easier to understand instructions from healthcare professionals³⁶, which aligns with our findings, where patients with partners exhibited better FHL. The association between marital status, being married or having a partner, and higher FHL may be related to the transfer of therapeutic responsibilities to the spouse.

FHL was not statistically associated with most clinical characteristics of patients, except for rehabilitation duration, which emerged as a significant predictor of adequate FHL. This finding corroborates other studies showing that clinical characteristics, such as injury level, were not associated with FHL^{17,24}.

Educating individuals with SCI to understand their health self-care needs is critical.

FUNCTIONAL HEALTH LITERACY IN INDIVIDUALS WITH SPINAL CORD INJURY TREATED AT A REHABILITATION CENTER

They require specialized attention and guidance to prevent complications and improve functionality. During the initial phase of physical rehabilitation, SCI patients may feel overwhelmed by their circumstances and struggle to process the large volume of information provided³⁶.

Longer rehabilitation durations increase exposure to health information, enhance organizational and management skills, and improve disease knowledge, positively impacting FHL outcomes. Clinical complications among SCI patients could be mitigated by assessing FHL levels prior to providing professional guidance, enabling personalized education programs to enhance rehabilitation outcomes and improve health results²⁴.

Although the FHL outcomes in SCI patients were satisfactory, this study has a limitation: it was conducted at a single rehabilitation center, which is a public facility specializing in SCI care. This context may limit the generalizability of the findings.

CONCLUSION

In the analysis of functional health literacy, the results were adequate for the majority of patients with spinal cord injury. In this study, younger age, higher income, greater educational attainment, longer rehabilitation duration, and having a partner were predictors of better FHL outcomes.

The quality of care provided, as well as the organizational structure of specialized and reference centers for SCI rehabilitation, can enhance outcomes related to FHL.

Identifying associated variables and analyzing the functional dimension of health literacy by multidisciplinary teams during rehabilitation are essential tools for improving patient health. However, it is important to acknowledge that assessing health literacy is complex due to its broad construct.

We recommend that future studies analyze the multidimensional aspects of health literacy in patients with SCI undergoing rehabilitation.

**FUNCTIONAL HEALTH LITERACY IN INDIVIDUALS WITH SPINAL
CORD INJURY TREATED AT A REHABILITATION CENTER**

REFERENCES

1. Badhiwala JH, Wilson JR, Fehlings MG. Global burden of traumatic brain and spinal cord injury. *Lancet Neurol.* 2019;18(1):24-25. DOI: [https://doi.org/10.1016/S1474-4422\(18\)30444-7](https://doi.org/10.1016/S1474-4422(18)30444-7)
2. Baldassin V, Shimizu HE. Como as tecnologias assistivas para acesso a computadores podem ajudar as pessoas com lesão medular? *RSD Journal* 2022; 11(2):971-979. DOI: <https://doi.org/10.33448/rsd-v11i2.19793>
3. Ding W, Hu S, Wang P, Kang H, Peng R, Dong Y, Li F. Spinal Cord Injury: The Global Incidence, Prevalence, and Disability From the Global Burden of Disease Study 2019. *Spine.* 2022;47(21):1532-1540. doi: : <https://doi.org/10.1097/BRS.0000000000004417>
4. Lu Y, Shang Z, Zhang W, Pang M, Hu X, Dai Y, Shen R, Wu Y, Liu C, Luo T, Wang X, Liu B, Zhang L, Rong L. Global incidence and characteristics of spinal cord injury since 2000-2021: a systematic review and meta-analysis. *BMC Med.* 2024;22(1):285. DOI: <https://doi.org/10.1186/s12916-024-03514-9>
5. Sobral Sousa S, Martins MM, Andrade MJ, Rodrigues Barbeiro S, Taveira Teixeira V. Cuidados de Enfermagem em Contexto Agudo à Pessoa com Lesão Medular: Scoping review. *Rev Port Enf Reab.* 2022;5(2):1-20. DOI: <https://doi.org/10.33194/rper.2022.204>
6. Singh G, Sawatzky B, Nimmon L, Mortenson WB. Perceived eHealth literacy and health literacy among people with spinal cord injury: A cross-sectional study. *J Spinal Cord Med.* 2023;46(1):118-125. doi: <https://doi.org/10.1080/10790268.2021.19631407>
7. Sørensen K, Van den Broucke S, Fullam J, Doyle G, Pelikan J, Slonska Z, Brand H. Health literacy and public health: A systematic review and integration of definitions and models. *BMC Public Health.* 2012; 12:80 DOI: <https://doi.org/10.1186/1471-2458-12-80>
8. Nutbeam D, Lloyd JE. Understanding and Responding to Health Literacy as a Social Determinant of Health. *Annu Rev Public Health.* 2021 Apr 1;42:159-173. DOI: <https://doi.org/10.1146/annurev-publhealth-090419-102529>
9. Sertkaya Z, Koyuncu E, Nakipoğlu, Yüzer GF, Özgirgin N. Investigation of health literacy level and its effect on quality of life in patients with spinal cord injury. *J Med Esp* 2021; 46:62-7. DOI: <https://doi.org/10.1080/10790268.2021.1991162>
10. Meggetto E, Kent F, Ward B, Keleher H. Factors influencing implementation of organizational health literacy: a realist review. *J Health Organ Manag.* 2020;34(5):325-407. DOI: <https://doi.org/10.1108/JHOM-06-2019-0167>
11. Gross-Hemmi MH, Gemperli A, Fekete C, Brach M, Schwegler U, Stucki G. Methodology and study population of the second Swiss national community survey of functioning after spinal cord injury. *Spinal Cord* 2021; 59(4):363-72. DOI: <https://doi.org/10.1038/s41393-020-00584-3>

**FUNCTIONAL HEALTH LITERACY IN INDIVIDUALS WITH SPINAL
CORD INJURY TREATED AT A REHABILITATION CENTER**

12. Parker RM, Ratzan S. Re-enforce, not re-define health literacy—moving forward with Health Literacy 2.0. *J Health Commun.* 2019;24(12):923-5. DOI: <https://doi.org/10.1080/10810730.2019.1691292>
13. Maragno, Carla Andreia Daros et al. Teste de letramento em saúde em português para adultos. *Rev Bras Epidemiol.* 2019;22: e190025. DOI: <https://doi.org/10.1590/1980-549720190025>.
14. Blacher PH, Mendonça IR, Vieira CS, Correa LK, Nunes FL, Martins LP et al. Relação entre o letramento em saúde e a adesão terapêutica de pacientes com diabetes mellitus tipo 2. *Arquivos Catarinenses de Medicina.* 2022; 51(3):124-134. Disponível em: <https://revista.acm.org.br/arquivos/article/view/1437>
15. Lima RIM, Parente MA, Ferreira TISP, Coelho AAS, Loureiro EVS de, Barbosa TM, Lustosa SB, Damasceno OC, Teixeira FB. Letramento funcional em saúde de usuários da atenção primária de Altamira, Pará: . *Rev Bras Med Fam Comunidade* . 2022;17(44):2763. DOI: [https://doi.org/10.5712/rbmfc17\(44\)2763](https://doi.org/10.5712/rbmfc17(44)2763)
16. Ribau A, Alves J, Rodrigues-Pinto R. Treatment of Acute Spinal Cord Injuries: A Survey Among Iberolatinoamerican Spine Surgeons - Part 2 Timing to Surgery. *Rev bras ortop* [Internet]. 2023Mar;58(2):337–41. DOI: <https://doi.org/10.1055/s-0042-1746181>
17. Lima AS, Lima BJS, Oliveira AT, Farias MGN, Passos MKA, Sandes MF, Santana LS, Rezende KF, Gomes ICP, Santana NO de. Health literacy in patients with chronic diseases. *RSD* [Internet]. 2022Jul.5 [cited 2024Oct.3];11(9): e15211931795-e15211931795. DOI: <https://doi.org/10.33448/rsd-v11i9.31795>
18. Chehuen JA, Costa LA, Estevanin GM, Bignoto TC, Vieira CIR, Pinto FAR, et al.. Letramento funcional em saúde nos portadores de doenças cardiovasculares crônicas. *Ciência saúde coletiva.* 2019;24(3):1121–32. DOI: <https://doi.org/10.1590/1413-81232018243.02212017>
19. Rodrigues da Silva JR, Andrade Luz GO de, da Silva SMB, Katiussia Araujo de Medeiros L, Santos Junior JL, Vieira Santos ICR. Letramento funcional em saúde e o conhecimento dos doentes renais crônicos em tratamento conservador. *Rev Bras Promoc Saúde.* 2019;32. DOI: <https://doi.org/10.5020/18061230.2019.9470>
20. Sutthiworapon S, Vichitkunakorn P, Choomalee K, Ngamchaliew P. Effect of online infographics for enhancing health literacy among patients with type 2 diabetes in primary care unit during the COVID-19 pandemic: a randomized controlled trial. *BMC Prim Care.* 2024 Mar 15;25(1):87. DOI: <https://doi.org/10.1186/s12875-024-02335-2>
21. Diviani N, Zanini C, Jaks R, Brach M, Gemperli A, Rubinelli S. Information seeking behavior and perceived health literacy of family caregivers of persons living with a chronic condition. The case of spinal cord injury in Switzerland. *Patient Educ Couns* 2020; 103:1531-7. <https://doi.org/10.1016/j.pec.2020.02.024>
22. Guan B, Anderson DB, Chen L, Feng S, Zhou H. Global, regional and national burden of traumatic brain injury and spinal cord injury, 1990-2019: a systematic analysis for the

**FUNCTIONAL HEALTH LITERACY IN INDIVIDUALS WITH SPINAL
CORD INJURY TREATED AT A REHABILITATION CENTER**

Global Burden of Disease Study 2019. *BMJ Open*. 2023;13(10):e075049. DOI: <https://doi.org/10.1136/bmjopen-2023-075049>

23. Silva HK de N da, Freitas C de A, Argenton VB, Calefi MPSS. Relação do tipo e nível de lesão medular espinhal com funcionalidade e qualidade de vida em um hospital de reabilitação. *REAS*. 2023;23(4):e12234. DOI: <https://doi.org/10.25248/reas.e12234.2023>

24. Silva FAR, Barbosa MA, Prudente COM, Morais LA, Moraes KL, Vila VSC, Porto CC. Health literacy of people with spinal cord injury: a systematic review. *Spinal Cord*. 2023 Aug;61(8):409-414. DOI: <https://doi.org/10.1038/s41393-023-00903-4>

25. Moriya KM, Condo TI, Montiel JM, Zanca GG. Health literacy and its relationship with quality of life, self-reported chronic diseases and difficulties for accessing health services among older adults. *RSD*. 2022;11(1):e3211124481. DOI: <https://doi.org/10.33448/rsd-v11i1.24481>

26. Pavão ALB, Werneck GL, Saboga-Nunes L, Sousa RA de. Avaliação da literacia para a saúde de pacientes portadores de diabetes acompanhados em um ambulatório público. *Cad Saúde Pública* 2021; 37(10):e00084819. DOI: <https://doi.org/10.1590/0102-311X00084819>

27. Campos AAL, Neves FS, Saldanha RF, Duque KCD, Guerra MR, Leite ICG, et al. Fatores associados ao letramento funcional em saúde de mulheres atendidas pela Estratégia de Saúde da Família. *Cad Saúde Colet*, 2020;28(1):66-76. DOI: <https://doi.org/10.1590/1414-462X202000280295>

28. Van Hoa H, Giang HT, Vu PT, Van Tuyen D, Khue PM. Factors Associated with Health Literacy among the Elderly People in Vietnam. *Biomed Res Int*. 2020 Mar 25;2020:3490635. DOI: <https://doi.org/10.1155/2020/3490635>

29. Chiu HT, Tsai HW, Kuo KN, Leung AYM, Chang YM, Lee PH, Hou WH. Exploring the Influencing Factors of Health Literacy among Older Adults: A Cross-Sectional Survey. *Medicina (Kaunas)*. 2020;56(7):330. DOI: <https://doi.org/10.3390/medicina56070330>

30. Abd-Rahim SNH, Mohamed-Yassin MS, Abdul-Razak S, Isa MR, Baharudin N. The Prevalence of Limited Health Literacy and Its Associated Factors among Elderly Patients Attending an Urban Academic Primary Care Clinic in Malaysia. *Int J Environ Res Public Health*. 2021;18(17):9044. DOI: <https://doi.org/10.3390/ijerph18179044>

31. Hochhauser M, Brusovansky M, Sirotin M, Bronfman K. Health literacy in an Israeli elderly population. *Isr J Health Policy Res*. 2019 Jul 10;8(1):61. DOI: 10.1186/s13584-019-0328-2. DOI: <https://doi.org/10.1186/s13584-019-0328-2>

32. Ma T, Meng H, Ye Z, Jia C, Sun M, Liu D. Health Literacy Mediates the Association Between Socioeconomic Status and Productive Aging Among Elderly Chinese Adults in a Newly Urbanized Community. *Front Public Health*. 2021 Apr 9;9:647230. doi: 10.3389/fpubh.2021.647230. DOI: <https://doi.org/10.3389/fpubh.2021.647230>

33. Soares TAM, Brasil VV, Moraes KL, Santos LTZ, Vila VSC, Borges Júnior LH. Health literacy of home caregivers in a Brazilian capital. *Acta Paul Enferm* 2021;34. DOI:

**FUNCTIONAL HEALTH LITERACY IN INDIVIDUALS WITH SPINAL
CORD INJURY TREATED AT A REHABILITATION CENTER**

<https://doi.org/10.37689/acta-ape/2021AO002255>

34. Buja A, Rabensteiner A, Sperotto M, Grotto G, Bertoncello C, Cocchio S, Baldovin T, Contu P, Lorini C, Baldo V. Health Literacy and Physical Activity: A Systematic Review. *J Phys Act Health*. 2020 Oct 31;17(12):1259-1274. DOI: <https://doi.org/10.1123/jpah.2020-0161>

35. Paes RG, Mantovani M de F, Costa MC, Pereira ACL, Kalinke LP, Moreira RC. Efeitos de intervenção educativa no letramento em saúde e no conhecimento sobre diabetes: estudo quase-experimental. *Esc Anna Nery* 2022; 26:e20210313. DOI: <https://doi.org/10.1590/2177-9465-EAN-2021-0313pt>

36. Stillman MD, Barber J, Burns S, Williams S, Hoffman JM. Complications of spinal cord injury over the first year after discharge from inpatient rehabilitation. *Archives of Physical Medicine and Rehabilitation* 2017; 98(9):1800-1805. DOI: <http://doi.org/0.1016/j.apmr.2016.12.011>

Submitted: October 31, 2023

Accepted: October 6, 2024

Published: March 17, 2025

Corresponding author:

Francine Aguilera Rodrigues da Silva: Conceptualization; Formal analysis; Investigation; Methodology, Project administration; Supervision; Validation; Visualization; Writing – original draft; Writing – review & editing.

Cejane Oliveira Martins: Supervision; Validation; Visualization; Writing – original draft; Writing – review & editing.

Vanessa da Silva Carvalho Vila: Visualization; Writing – original draft; Writing – review & editing.

Katarinne Lima Moraes: Visualization; Writing – original draft; Writing – review & editing.

Maria Alves Barbosa: Project administration; Supervision; Validation; Visualization; Writing – original draft; Writing – review & editing.

Celmo Celeno Porto: Supervision; Validation; Supervision; Visualization; Writing – original draft; Writing – review & editing.

All authors approved the final version of the text.

Conflict of interest: There is no conflict of interest.

**FUNCTIONAL HEALTH LITERACY IN INDIVIDUALS WITH SPINAL
CORD INJURY TREATED AT A REHABILITATION CENTER**

Funding: Programa de Pós-graduação em Ciências da Saúde da Universidade Federal de Goiás

Corresponding author:

Francine Aguilera Rodrigues da Silva

Universidade Federal de Goiás – UFG

Rua 235. s/n. Setor Leste Universitário. CEP.74605-050. Goiânia/GO, Brasil.

francine.fisio@hotmail.com

Editor-in-chief: Adriane Cristina Bernat Kolankiewicz. PhD

This is an open access article distributed under the terms of the Creative Commons license.

