

EVALUATION OF THE WORK PROCESS OF PRIMARY HEALTH CARE TEAMS IN PRENATAL CARE

Érica de Brito Pitilin¹; Daniela Savi Geremia²; Jéssica Ferreira³
Yaná Tamara Tomasi⁴; Carine Vendruscolo⁵; Tatiane Baratieri⁶
Maicon Henrique Lentsck⁷; Janine Schirmer⁸

Highlights: (1) Managerial and care factors influence access to prenatal consultations. (2) Poor referral flow and communication hinder effective prenatal care. (3) Continuing education and care safety promote prenatal consultations.

PRE-PROOF

(as accepted)

This is a preliminary, unedited version of a manuscript accepted for publication in Revista Contexto & Saúde. As a service to our readers, we are making this initial version of the manuscript available, as accepted. The article will still be reviewed, formatted, and approved by the authors before being published in its final form.

<http://dx.doi.org/10.21527/2176-7114.2025.50.14540>

How to cite:

Pitilin É de B, Geremia DS, Ferreira J, Tomasi YT, Vendruscolo C, Baratieri T. et al. Evaluation of the work process of primary health care teams in prenatal care. Rev. Contexto & Saúde, 2025;25(50):e14540

¹ Universidade Federal da Fronteira Sul. Chapecó/SC, Brasil. <https://orcid.org/0000-0003-3950-2633>

² Universidade Federal da Fronteira Sul. Chapecó/SC, Brasil. <https://orcid.org/0000-0003-2259-7429>

³ Universidade Federal de Minas Gerais. Belo Horizonte/MG, Brasil. <https://orcid.org/0000-0002-8284-7883>

⁴ Universidade Federal da Fronteira Sul. Passo Fundo/RS, Brasil. <https://orcid.org/0000-0001-6034-1497>

⁵ Universidade do Estado de Santa Catarina. Chapecó/SC, Brasil. <https://orcid.org/0000-0002-5163-4789>

⁶ Universidade Estadual do Centro-Oeste. Guarapuava/PR, Brasil. <https://orcid.org/0000-0002-0270-6395>

⁷ Universidade Estadual do Centro-Oeste. Guarapuava/PR, Brasil. <https://orcid.org/0000-0002-8912-8902>

⁸ Universidade Federal de São Paulo. São Paulo/SP, Brasil. <https://orcid.org/0000-0003-0783-2961>

EVALUATION OF THE WORK PROCESS OF PRIMARY HEALTH CARE TEAMS IN PRENATAL CARE

ABSTRACT

The objective was to identify the factors associated with prenatal care based on the work process of primary health care teams in Brazil. This is a cross-sectional study with secondary data from 38,865 teams interviewed who took part in the external evaluation of the third cycle of the National Program for Improving Primary Care Access and Quality (PMAQ-AB, acronym in Portuguese) in 2017. The dependent variable was prenatal care, and the independent variables were grouped into hierarchical levels. Multiple logistic regression analysis was conducted, and the variables were classified into management and care. The care factors associated with consultation included continuing education, safety in providing care, and enough professionals. An institutionalized communication flow with the specialist team is absent in the managerial dimension. The lack of defined flows for delivery/maternity and the absence of a central regulation system for referrals, among others, contributed to the pregnant woman not having a consultation. The analysis provides input for improving work processes in prenatal care and strengthening the integration of the care network.

Keywords: Primary Health Care; Prenatal Care; Health Evaluation.

INTRODUCTION

Prenatal care (PNC) is one of the services that pregnant women need to follow up on and monitor the progress of their pregnancy, as it is one of the leading indicators of the proposed United Nations (UN) Sustainable Development Goals¹. Recently, several initiatives have been implemented to improve the quality of care for pregnant women in Primary Health Care (PHC). The National Program for Improving Primary Health Care Access and Quality (PMAQ-AB) is one of the most relevant health strategies and the most significant proposal for evaluating the performance of programs ever developed in Brazil^{2,3}. This program aims to encourage managers and teams to improve the quality of services through strategies for qualifying, monitoring, and evaluating the work of health teams⁴. Although the PMAQ still has limitations, its potential is represented by the participatory process of the entire team in the evaluation processes. In Brazil, PHC is referred to as Primary Care.

EVALUATION OF THE WORK PROCESS OF PRIMARY HEALTH CARE TEAMS IN PRENATAL CARE

Despite recent advances in social and demographic indicators in Brazil, inequalities persist in access to and quality of PNC, especially in less developed regions⁵. These inequalities can be seen in the access, declaration, and outcome of PNC⁶. In this sense, Primary Care has expanded its offer of consultations and procedures to develop resolute and welcoming actions with the health services network. However, there is a lack of evaluation studies, monitoring indicators, and management actions that favor the quality of the PNC offered by teams⁷.

In several countries, problems relating to the quality of PNC have also been highlighted. In the maternal care systems of Central and Eastern Europe, inequalities persist with barriers to access and maternal care, as well as inadequate and outdated protocols, even though these systems are well implemented on the continent⁸.

A study in some regions of Asia identified that the quality of PNC is inadequate, precisely due to managerial and care factors, such as the availability of human resources, facilities, distance between units, and more⁹. Another study, conducted in a Latin American context, identified low adherence to prenatal care and difficulties for health teams to link up with other points in the health care network¹⁰.

From this perspective, evaluating the completeness of a repertoire of health actions is valuable for estimating the scope of care for pregnant women and its quality¹¹. Information on the work processes of health teams and the organization of care for the user can support management and care strategies, which will help evaluate the quality of prenatal care in the country. This study aims to identify the factors associated with prenatal care based on the work process of PHC teams in Brazil.

METHODS

This cross-sectional study uses secondary data from PHC teams in the Brazilian Unified Health System (SUS), whose professionals responded to the external evaluation component of the third cycle of the PMAQ-AB. This program is linked to financial transfers based on the certification result obtained by the teams, which covered 93.9% of PHC teams in 95.6% of the municipalities in Brazil in 2017 when the data was collected¹².

EVALUATION OF THE WORK PROCESS OF PRIMARY HEALTH CARE TEAMS IN PRENATAL CARE

The PMAQ-AB had an evaluation tool comprising three modules that guided data collection. Module I described the observation of the structural characteristics and ambience of the health unit; Module II interviewed professionals about the health team's work process and analyzed documents; and Module III interviewed users about their perceptions and satisfaction with health services in terms of access and use¹². For this study, the variables contained in Module II were used so that the interviews took place using a non-probabilistic sample, in which four users from each unit were interviewed, totaling 140,444 users nationwide¹².

PMAQ data is collected through a tablet application, where all those involved in the health teams actively collaborate to ensure the smooth running of the program's external evaluation phase. The Module II variables were accessed by consulting the PMAQ *website* (<https://aps.saude.gov.br/ape/pmaq>), available as an XLSX file¹².

This study's population included teams that answered “yes” to “Do you provide prenatal care?”, resulting in 38,865 questionnaires.

Whether or not prenatal consultations were conducted was considered a dependent variable, and the determinants of whether consultations were undertaken in PHC were classified into two dimensions of analysis: management and care. The conceptual model was based on Donabedian's theoretical framework¹³ and adapted to represent the components of prenatal care (APN) proposed by Luz, Aquino, and Medina¹⁴.

The components of the PNC and their relationships with each other were formulated in two dimensions of analysis: distal management level and proximal care level, made up of structure and process attributes (Figure 1).

The analysis matrix for the distal management level comprises structure and process attributes and includes the following variables: Is there an institutionalized communication flow between your team and specialist care? Do specialists contact primary care professionals to exchange information about pregnant women referred by primary care? Are there defined referrals and flows for delivery (maternity)? Is a regulation center available to refer pregnant women to other points of care? In the last year, has the team taken part in continuing education? Do the continuing educational activities meet the demands and needs of the team? Does the team monitor and analyze health indicators and information? Is there a lack of training?

EVALUATION OF THE WORK PROCESS OF PRIMARY HEALTH CARE TEAMS IN PRENATAL CARE

In the care dimension, the structure and process variables were considered: Does the team receive support from other professionals to help or support the resolution of complex cases? Support from the Expanded Family Health and Primary Care Centers (NASF-AB)? Do you know if the team conducts any activities to plan its actions? Is the infrastructure inadequate? Weren't the materials/supplies enough? Is there an absence of parameters? Is there a deficit of professionals (human resources)? Is there a lack of demand for the service? Work overload? Are there restrictive measures by the profession's supervisory board? Is there a lack of security during the consultation?

Most of the time, each criterion was constructed by aggregating aspects from more than one question from the PMAQ-AB instruments, dichotomizing the answer alternatives into yes and no. For the questions relating to actions taken by health professionals, only those cases in which the action was reported with documentary confirmation were considered to have an affirmative answer. Finally, the answer options 'never available/sometimes available', 'don't know/didn't answer', and 'not applicable' were classified as 'no'.


After selecting the variables, the descriptive analysis of the database identified 23,067 interviews with missing answers (*missing*) in at least one variable. The multiple imputation method was used to avoid excluding participants, where missing data is imputed **m** times, generating **m** distinct and complete databases. From these different databases, we obtain results that match the desired objective, considering the variability of the imputation¹⁵.

The imputation model is specified separately for each variable, using all the other variables as predictors. At each algorithm stage, an imputation is generated for the missing data variable; these imputed values are used to impute the following variable. The process continues until convergence is reached. In other words, the basic idea is to impute the incomplete variables one at a time, using all the variables and the variables already imputed in the previous stage.^{16,17}

Multiple logistic regression with hierarchical variable entry was used for the analysis. Initially, the distal variables, considered conditioning factors for the variables at the next level, were analyzed; then, the intermediate and proximal variables were examined.

EVALUATION OF THE WORK PROCESS OF PRIMARY HEALTH CARE TEAMS IN PRENATAL CARE

Figure 1 - Subdimensions of the National Program for Improving Primary Care Access and Quality database, 2017.



Source: PMAQ-AB Adaptation, 2017.

The multiple logistic regression model, including *stepwise forward* variables, considered those with a p-value < 0.20 in the bivariate analysis, and the variables with a p-value < 0.05 and/or which adjusted the model remained in the final model. The magnitude of the associations was estimated by the *Odds Ratio* (OR), with 95% confidence intervals (95%CI) as a measure of precision. The adequacy of the final model was checked using the Hosmer-Lemeshow test. The multicollinearity assumption was tested using the *Variance Inflation Factor* (VIF) and calculated by multiple linear regression, with the cut-off point being $VIF \geq 4$. The test showed no evidence of multicollinearity between the independent variables studied for multiple logistic regression. The data was stored and analyzed using the *Statistical Package for the Social Sciences* (SPSS) *software*, version 20.0.

As this study uses secondary databases, whose information is aggregated without the possibility of individual identification, the Human Research Ethics Committee is exempt from analyzing it under Resolution 510/2016-of the National Health Council.

RESULTS

The proportion of teams that carried out prenatal consultations was 96.9% (n=36,702), with 35.6% (n=13,486) working in the northeast, followed by 32.9% (n=12,482) in the southeast and 14.7% (n=5,558) in the south. Table 1 shows the descriptive analysis of the primary care teams

EVALUATION OF THE WORK PROCESS OF PRIMARY HEALTH CARE TEAMS IN PRENATAL CARE

in terms of their employment characteristics. There was a predominance of nurses, 93.3% (n=35,371), who were civil servants and had a career plan.

Table 1 - Characteristics of health professionals in terms of employment profile, 3rd cycle of the PMAQ-AB. Brazil, 2017.

Variables	Population analyzed	n 37,894	% 100
Region	North	3,714	9.8
	Northeast	13,486	35.6
	South	5,558	14.7
	Southeast	12,482	32.9
	Midwest	2,654	7.0
Professional interviewed	Nurse	35,371	93.3
	Doctor	1,869	4.9
	Other senior professionals	110	0.3
What type of link	Civil servant	16,054	42.4
	Commissioned position	1,078	2.8
	Temporary contract by the public administration	8,530	22.5
	Temporary public administration contracts	4,726	12.5
	CLT public employment	2,321	6.1
	CLT Contract	3,992	10.5
	Self-employed	107	0.3
	Other	542	1.4
Career plan	Seniority progression	8,302	21.9
	Progression for performance/merit	5,674	15.0
	Progression for qualifications and training	8,144	21.5

Source: Authors, 2023.

Regarding the variables related to the work process at the distal level, on the anagerial dimension, the bivariate analysis indicated that all the variables were significant in at least one of their categories, as shown in Table 2.

EVALUATION OF THE WORK PROCESS OF PRIMARY HEALTH CARE TEAMS IN PRENATAL CARE

Table 2 - Bivariate analysis of the association between the variables related to the distal level in the management dimension by aggregating aspects from the PMAQ-AB and the prenatal visit. Brazil, 2017.

Variables	Conducts prenatal consultations		OR _{crude} (CI95%)*	p-value
	Yes n (%) 36,702 (96.9)	No n (%) 1,192 (3.1)		
Is there an institutionalized communication flow between your team and specialized care?				
Yes	32,835 (87.9)	548 (1.5)	1.00	Reference
No	3,867 (10.4)	100 (0.3)	1.54 (1.24 – 1.92)	0.000
Do the specialists get in touch to exchange information about pregnant women referred by AB?				
Yes	26,074 (78.1)	399 (1.2)	1.00	Reference
No	6,761 (20.3)	149 (0.4)	1.44 (1.19 – 1.74)	0.000
There are defined referrals and flows for childbirth (maternity).				
Yes	31,425 (94.1)	493 (1.5)	1.00	Reference
No	1,410 (4.2)	55 (0.2)	2.48 (1.87- 3.30)	0.000
Is a regulation center available to refer pregnant women to other points of care?				
Yes	34,844 (93.3)	598 (1.6)	1.00	Reference
No	1,858 (5.0)	50 (0.1)	1.56 (1.17 – 2.10)	0.000
In the last year, has the team taken part in continuing education?				
Yes	35,668 (95.5)	600 (1.6)	0.36 (0.26 – 0.48)	0.000
No	1,034 (2.8)	49 (0.1)	1.00	Reference
Do continuing education actions meet the demands and needs of the team?				
Yes	35,371 (97.5)	587 (1.6)	1.00	Reference
No	297 (0.8)	13 (0.0)	2.63 (1.50 – 4.62)	
Does the team monitor and analyze health indicators and information?				
Yes	32,375 (86.7)	468 (1.3)	1.00	Reference
No	4,327 (11.6)	180 (0.5)	2.87 (2.41 – 3.42)	0.000
There is a lack of training.				
Yes	2,553 (6.9)	37 (0.1)	1.20 (0.86 – 1.67)	0.160
No	33,948 (91.4)	591 (1.6)	1.00	Reference

*95% CI: 95% confidence interval; OR: odds ratio - crude value.

Source: Authors, 2023.

Of the proximal level variables related to the care dimension included in the bivariate analysis, those significantly affecting the study outcome were support from other professionals in the care network in resolving complex cases, support from the NASF-AB, health action planning activities, available human resources, and safety in providing care to pregnant women (Table 3).

EVALUATION OF THE WORK PROCESS OF PRIMARY HEALTH CARE TEAMS IN PRENATAL CARE

Table 3 - Bivariate analysis of the association between the variables related to the proximal level of care by aggregating aspects from the PMAQ-AB and the prenatal visit. Brazil, 2017.

Variables	Conducts prenatal consultations		OR _{crude} CI(95%)*	p-value
	Yes n (%) 36,702 (96,9)	No n (%) 1,192 (3,1)		
<i>Does the team receive support from other professionals when solving complex cases?</i>				
Yes	35,923 (96.2)	624 (1.7)	1.00	Reference
No	779 (2.1)	24 (0.1)	1.77 (1.17 – 2.68)	0.008
<i>Is there NASF-AB support?</i>				
Yes	26,887 (73.6)	326 (0.8)	1.00	Reference
No	9,036 (24.7)	298 (0.8)	2.70 (2.32 – 3.18)	0.000
<i>Does the team carry out any activities to plan its actions?</i>				
Yes	35,478 (95.0)	549 (1.5)	1.00	Reference
No	1,224 (3.3)	99 (0.3)	5.22 (4.18 – 6.52)	0.000
<i>Is there inadequate infrastructure?</i>				
Yes	7,785 (21.0)	140 (0.4)	1.00	Reference
No	29,716 (77.3)	488 (1.3)	0.94 (0.78 – 1.14)	0.296
<i>Is there a shortage of materials?</i>				
Yes	7,961 (21.4)	138 (0.4)	1.00	Reference
No	28,540 (76.9)	490 (1.3)	0.99 (0.81 – 1.19)	0.480
<i>Is there a lack of protocols?</i>				
Yes	4,539 (12.2)	81 (0.2)	1.00	Reference
No	31,962 (86.1)	547 (1.5)	0.95 (0.75 – 1.21)	0.387
<i>Is there a shortage of professionals (human resources)?</i>				
Yes	2,406 (6.5)	52 (0.1)	1.00	Reference
No	34,095 (91.8)	576 (1.6)	0.78 (0.58 – 1.04)	0.054
<i>Is there a lack of demand for the service?</i>				
Yes	5,295 (14.3)	91 (0.2)	1.00	Reference
No	31,206 (84.0)	537 (1.4)	1.00 (0.80 – 1.25)	0.518
<i>Is there a work overload?</i>				
Yes	1,865 (5.0)	29 (0.1)	1.11 (0.76 – 1.61)	0.321
No	34,636 (93.3)	599 (1.6)	1.00	Reference
<i>Are there any restrictive measures from the profession's supervisory board?</i>				
Yes	1,092 (2.9)	22 (0.1)	0.84 (0.55 – 1.30)	0.265
No	35,409 (95.4)	606 (1.6)	1.00	Reference
<i>Is there a lack of security when providing care to pregnant women?</i>				
Yes	1,622 (4.4)	35 (0.1)	0.78 (0.55 – 1.11)	0.104
No	34,978 (93.9)	593 (1.6)	1.00	Reference

*95% CI: 95% confidence interval; OR: odds ratio - crude value.

Source: Authors, 2023.

Table 4 shows the hierarchical multiple analysis models by aggregating aspects from the PMAQ-AB and prenatal care. Model A, referring to the distal-level variables, and Model

EVALUATION OF THE WORK PROCESS OF PRIMARY HEALTH CARE TEAMS IN PRENATAL CARE

B, referring to the proximal-level variables, include all the statistically significant variables in the multiple analysis.

The multiple analysis showed that factors such as the absence of an institutionalized communication flow with the specialist team, as well as contact between specialists and PHC professionals about pregnant women, the absence of defined flows for childbirth/maternity, the absence of a regulation center available for referrals to other points of care, the lack of monitoring and analysis of health indicators and information, and the absence of capacity building/training were more likely to cause teams not to carry out prenatal consultations. It should be noted that the lack of planning of the team's actions was 2.95 times more likely not to provide care to pregnant women (OR 2.95, 95% CI 2.67 - 3.26) (Table 4).

On the other hand, factors such as continuing education, sufficient professionals/human resources, and safety in providing care to pregnant women functioned as favorable factors in providing prenatal care (Table 4).

Table 4 - Hierarchical multiple logistic regression analysis aggregates aspects from the PMAQ-AB and prenatal care. Brazil, 2017.

Variables	Model A* Adjusted OR(95%CI)***	p-value	Model B** Adjusted OR(95%CI)	p-value
Lack of institutionalized communication flow with the specialist team.	1.18 (1.07 – 1.30)	0.001	1.10 (1.00 – 1.20)	0.034
Lack of contact between specialists and primary care professionals about pregnant women.	1.15 (1.07 – 1.23)	0.000	1.08 (1.01 – 1.16)	0.025
Lack of defined delivery/maternity flows.	1.83 (1.64 – 2.05)	0.000	1.83 (1.64 – 2.05)	0.000
No central regulation center available for referrals to other points of care.	1.16 (1.03 – 1.31)	0.011	1.15 (1.02 – 1.30)	0.019
Continuing education.	0.54 (0.47 – 0.62)	0.000	0.66 (0.58 – 0.76)	0.000
Continuing educational actions meet the demands and needs of the team.	1.69 (1.34 – 2.14)	0.000	0.41 (0.12 – 1.79)	0.304
Failure to monitor and analyze health indicators and information.	2.28 (2.12 – 2.46)	0.000	1.75 (1.62 – 1.90)	0.000
Lack of training.	1.15 (1.01 – 1.30)	0.030	1.45 (1.25 – 1.68)	0.000
Lack of NASF-AB support.	-	-	2.16 (2.03 – 2.30)	0.000
Lack of action planning by the AB team.	-	-	2.95 (2.67 – 3.26)	0.000
Enough professionals/human resources.	-	-	0.78 (0.69 – 0.88)	0.000
Safety in providing care to pregnant women.	-	-	0.70 (0.60 – 0.82)	0.000

* Model of the variables related to the distal level in the management dimension.

** Model of the variables related to the proximal level of the care dimension adjusted by the distal level variables.

***IC95%: 95% confidence interval; OR: odds ratio - adjusted value.

Source: Authors, 2023.

EVALUATION OF THE WORK PROCESS OF PRIMARY HEALTH CARE TEAMS IN PRENATAL CARE

DISCUSSION

This study demonstrated essential factors in the work process of PHC health teams, both in terms of management and care and whether prenatal consultations are carried out in the country. In Brazil, most studies related to prenatal care have focused their analysis on classic user evaluation criteria^{2,6,18}.

The use of PMAQ-AB data has strengthened the evaluation process in the SUS and revealed, considering Donabedian's¹³ approach, the management conditions and organization of the work process of health unit professionals and the development of health actions within the scope of PHC¹⁴. From a theoretical and methodological point of view, PMAQ-AB has broadened the scope of analysis of PNC in the country with the availability of indicators and information that cover issues of access to maternal care outcomes.

Despite the high coverage of PNC and its institutionalization many years ago in standards and protocols in Brazil, problems of various managerial natures and working conditions persist, potentially damaging maternal and child health^{2,6}.

It is worth noting that the difficulty in developing management actions in health units and managing policies at the municipal level are challenges for improving PNC throughout the country. As for the management conditions that have the potential to support care actions, this study identified difficulties in planning and organizing flows in the network for monitoring pregnant women, a lack of regulation, and a deficit in the use of indicators, monitoring, and analysis of data that support activities related to PNC.

Furthermore, this study corroborates the results found in another evaluation of prenatal care at a national level⁷ regarding the inadequacy of the (infrastructure) structure of basic units for maternal care, which affected approximately 43% of the units surveyed in the study. This is pointed out as a barrier to the quality of PHC in local scenarios in country¹⁹.

In addition to the managerial dimension, investigating the determinants of why professionals in the teams do not carry out prenatal consultations can have a negative influence on the longitudinality of care, on the team's relationships with the populations they assist, on pregnant women's adherence to prenatal care, and, consequently, on the inefficiency of PHC²⁰. The principle of continuous care implies trust between user and professional, as it implies trust between user and professional. This attribute of PHC is of great importance in prenatal care so

EVALUATION OF THE WORK PROCESS OF PRIMARY HEALTH CARE TEAMS IN PRENATAL CARE

that there is continuity of care for pregnant women, ensuring the bond between women and health professionals to achieve a good progression of pregnancy and the well-being of pregnant women²¹. It is, therefore, essential to guarantee the availability of consultations and the monitoring of pregnancy progress for all pregnant women in all units across the country.

Studies have assessed the quality of prenatal care and identified weaknesses in care when teams do not have matrix support from other professionals and/or other points in the healthcare network. They have also identified a lack of planning and organization of the work process, well-defined flows, and guidelines.^{18,22,23}

This study's lack of support meant that PHC team professionals were twice as likely not to conduct prenatal consultations. Matrix support has helped to improve the quality of PHC in Brazil, and important levels of matrix support corresponded to high chances of teams obtaining better certification in women's care by the PMAQ-AB²⁴.

The idea that characterizes matrix support is linked to that of the reference team - a discussion presented by Campos (1999)²⁵ - and is a different operating mode from the traditional one, in which “specialized” professionals are responsible for providing pedagogical support to the professionals in the generalist PHC teams²⁵. The results of the NASF-AB work process represented a fascinating evolution in the work of PHC teams in terms of collaborative action²⁶.

However, the change in the National Primary Care Policy on September 21, 2017, in the organization and funding for the Nuclei led to their extinction in many places, not guaranteeing that the logic of matrix support would remain, even in the territories in which the “support” multi-professional teams persist²⁷.

Other factors, such as the lack of care flows, available central regulation, and monitoring and analysis of health information indicators, were also crucial in the PHC teams' failure to conduct prenatal consultations. These aspects are essential for establishing the basis for the referral system for the country's pregnant women, which leads to discontinuity and fragmentation of care²⁸. This implies the need to continue discussing the attributes of PHC in terms of the coordination of care (in this case, the lack of it).

The definition of care coordination was primarily inspired by the conceptual model that accurately illustrates the meaning attributed to care coordination: *anything that bridges gaps*.

EVALUATION OF THE WORK PROCESS OF PRIMARY HEALTH CARE TEAMS IN PRENATAL CARE

In this sense, coordination means establishing connections to meet users' needs and preferences in providing quality care. In PHC, the absence of roles and flows aggravates this disarticulation of care and assistance, implying the quality of regulation processes, including the insufficient supply of specialized care in the SUS²⁹.

The network fragmentation, lack of flow definition, and disruption of health services make it even more difficult for the PNC to perform its role as the preferred point of entry for the care of pregnant women. Each point of care in the network must be appropriately structured so that the work process can be organized and at least one line of care is followed for the demands of pregnant women.

Despite the guidelines for achieving comprehensive prenatal care (PNC), such as the Stork Network and the Prenatal and Birth Humanization Policy (PHPN), it is necessary to rethink how these policies are implemented and to rethink prenatal care models based on the uniqueness of each woman, family, and territory³⁰.

In this context, the organization of the work process is linked to the quality of prenatal care to guarantee the sustainability of actions and the agility of the maternal care flow (management dimension vs. care dimension). Even though most of the teams interviewed said they had well-defined care protocols and flows within the care network, the need to guarantee and provide comprehensive and resolute care for all pregnant women in the country is reinforced.

The regulation of pregnant women's care is the responsibility of managers, permeated by public, private, and individual motivations and interests. However, it impacts the work process of the PHC team. Therefore, managers should provide spaces for dialogue and joint planning between professionals in the service network and the regulation centers, with a view to systemic integration and defragmentation of the care flow.

On the other hand, factors such as continuing education, safety in providing care to pregnant women, and enough professionals/human resources reinforced the prenatal care provided by the PHC teams in this study. Therefore, this is a significant health care initiative (including prenatal care) and should be instituted regularly in institutions.

Investing in health professionals' qualifications strengthens their capacity for planning, collective, and collaborative work. It also enhances cognitive tools such as welcoming,

EVALUATION OF THE WORK PROCESS OF PRIMARY HEALTH CARE TEAMS IN PRENATAL CARE

guaranteeing agile flow, and coordination between the different points of health care, which are fundamental for conducting prenatal consultations. In addition, enough professionals in the team's enhance these tools' use, favoring the professionals' work process and guaranteeing comprehensive and coordinated care for pregnant women ²⁶.

It highlights the critical role that primary care (PC) teams play in caring for and monitoring pregnant women in Brazil and their direct relationship with better morbidity and mortality indicators.

CONCLUSION

The work process of the PHC teams in prenatal care was associated with both management and care factors. The percentage of teams identified in this study with resources and strategies available to conduct prenatal care in the units was quite significant. However, ensuring that all pregnant women in the country have access to prenatal care and monitor their pregnancy progress is essential.

Factors such as continuing education, safety in providing care to pregnant women, and sufficient professionals/human resources functioned as factors within the care dimension that favor and reinforce the provision of prenatal care by health teams. On the other hand, factors in the management dimension such as the absence of an institutionalized communication flow with the specialist team, the lack of contact between specialists and PHC professionals about pregnant women, the absence of defined flows for delivery/maternity, the absence of a central regulation system available for referrals to other points of care, among others, were factors that contributed to the teams analyzed not carrying out the consultation for pregnant women.

The PMAQ-AB proved to be a valuable tool for evaluating most of the items relating to health professionals' work process in Brazil's primary care teams. This instrument was developed to assess Primary Care (AB, in Portuguese) and not exclusively evaluate prenatal care. It was possible to extract extensive elements relating to management and organization to analyze prenatal care. However, the evaluation through the PMAQ-AB is carried out because of adherence by the teams, i.e., it does not represent all of them. It may have a selection bias because more responses were concentrated in some regions than others, which may be a study limitation.

EVALUATION OF THE WORK PROCESS OF PRIMARY HEALTH CARE TEAMS IN PRENATAL CARE

The innovative nature of the work process analysis using PMAQ-AB indicators concerning maternal health stands out for identifying positive and negative factors in management conditions and the organization (or lack of) of teams in planning care that allows users to follow a therapeutic path or flow of care, supported by a line of care and guaranteeing quality access. It thus provides input for rethinking work processes and strengthening the integration of the care network based on initiatives between managers, regulation centers, and health teams.

Based on subsequent external evaluations of the PMAQ-AB, other studies should be conducted to assess the quality of care provided to pregnant women in the country and observe adjustments over time.

REFERENCES

1. Santos LKR, Oliveira F, Bastos JL. Iniquidades da assistência pré-natal no Brasil: uma análise interseccional. *Physis*. 2024;34:e34004. DOI: <https://doi.org/10.1590/S0103-7331202434004pt>
2. Medeiros GAR, Nickel DA, Calvo MCM. Construindo um modelo para avaliar o uso do Programa Nacional de Melhoria do Acesso e da Qualidade da Atenção Básica. *Epidemiol Serv Saude*. 2019;28(3):e2018281. DOI: <https://doi.org/10.5123/S1679-49742019000300006>
3. Brasil. Ministério da Saúde (MS). Saúde mais perto de você – acesso e qualidade. Programa nacional de melhoria do acesso e da qualidade da atenção Básica (PMAQ-AB): documento síntese para avaliação externa. Brasília: Ministério da Saúde; 2012.
4. Brasil. Ministério da Saúde (MS). Portaria MS/GM nº. 1.645, de 02 de outubro de 2015. Dispõe sobre o Programa Nacional de Melhoria do Acesso e da Qualidade da Atenção Básica. *Diário Oficial da União*; 2 out 2015 [acessado 06 fev 2024]. Disponível em: https://bvsms.saude.gov.br/bvs/saudelegis/gm/2015/prt1645_01_10_2015.html
5. Russo LX, Powell-Jackson T, Maia Barreto JO, Borghi J, Kovacs R, Gurgel Junior GD, et al. Pay for performance in primary care: the contribution of the Programme for Improving Access and Quality of Primary Care (PMAQ) on avoidable hospitalisations in Brazil, 2009 – 2018. *BMJ Global Health*. 2021;6(7):e005429. DOI: <https://doi.org/10.1136/bmjgh-2021-005429>
6. Rosa VM, Roxana Knobel, Traebert ESA, Iser BPM. Assessment of prenatal care adequacy using different normative criteria in a municipality in Santa Catarina, Brazil. *International Journal of Population Studies*. 2024;10(3):17–33. DOI: <https://doi.org/10.36922/ijps.1422>

**EVALUATION OF THE WORK PROCESS OF
PRIMARY HEALTH CARE TEAMS IN PRENATAL CARE**

7. Silva JP, Fernandes Costa D. O impacto dos fatores socioeconômicos na qualidade da assistência do pré-natal na atenção primária no Brasil. *Rev CEREUS*. 2024;16(2):333-351. DOI: <https://doi.org/10.18605/2175-7275/cereus.v16n2p333-351>
8. Lunda P, Minnie CS, Lubbe W. Factors influencing respectful perinatal care among healthcare professionals in low-and middle-resource countries: a systematic review. *BMC Pregnancy Childbirth*. 2024;442. DOI: <https://doi.org/10.1186/s12884-024-06625-6>
9. Sarikhani Y, Najibi SM, Razavi Z. Key barriers to the provision and utilization of maternal health services in low-and lower-middle-income countries; a scoping review. *BMC Women's Health*. 2024;325. DOI: <https://doi.org/10.1186/s12905-024-03177-x>
10. Simarmata M. Basic Maternal Care for Pregnant Women Under The Regulation of Minister of Health Number 4 of 2019. *Proceedings: International Forum Research on Education, Social Sciences Technology and Humanities*. 2024;1(2):118–127. Retrieved from <https://journal.berpusi.co.id/index.php/POE/article/view/850>
11. Nahidi F, Hajifoghaha M, Simba M, Nasiri M. Assessment of Prenatal Care Providers' Competencies From the Perspective of Pregnant Women: An Iranian Study. *J Patient Exp*. 2022;9:1-7. DOI: <https://doi.org/10.1177/23743735221092559>
12. Brasil. Secretaria de Atenção Primária à Saúde. Ministério da Saúde (MS). Programa Nacional de Melhoria do Acesso e da Qualidade da Atenção Básica (PMAQ) [site]. 2017 [acessado em 23 nov 2020]. Disponível em: <https://aps.saude.gov.br/ape/pmaq>
13. Donabedian A. Garantía y Monitoría de Calidad de la Atención Médica: Un texto introductorio. México: Instituto Nacional de Salud Pública; 1990.
14. Luz LA, Aquino R, Medina MG. Avaliação da qualidade da Atenção Pré-Natal no Brasil. *Saude Debate*. 2018;42(2):111-126. DOI: <https://doi.org/10.1590/0103-11042018S208>
15. Nunes LN, Klück MM, Fachel JMG. Comparação de métodos de imputação única e múltipla usando como exemplo um modelo de risco para mortalidade cirúrgica. *Rev bras epidemiol*. 2010;13(4):596-606. DOI: <https://doi.org/10.1590/S1415-790X2010000400005>
16. Horton NJ, Kleinman KP. Much ado about nothing: A comparison of missing data methods and software to fit incomplete data regression models. *Am Stat*. 2007;61(1):79-90. DOI: <https://doi.org/10.1198/000313007X172556>
17. Dong Y, Chao-Ying JP. Principled missing data methods for researchers, SpringerPlus 2013;2(1):222. DOI: <https://doi.org/10.1186/2193-1801-2-222>
18. Fausto MCR, Giovanella L, Mendonça MHM de, Seidl H, Gagno J. A posição da Estratégia Saúde da Família na rede de atenção à saúde na perspectiva das equipes e usuários participantes do PMAQ-AB. *Saude debate*. 2014;38(esp):13-33. DOI: <https://doi.org/10.5935/0103-1104.2014S003>

**EVALUATION OF THE WORK PROCESS OF
PRIMARY HEALTH CARE TEAMS IN PRENATAL CARE**

19. Ferreira J, Geremia DS, Geremia F, Celuppi IC, Thomas TLH, Souza JB. Avaliação da Estratégia Saúde da Família à luz da tríade de Donabedian. *Av enferm.* 2021;39(1): 63-73. DOI: <https://doi.org/10.15446/av.enferm.v39n1.85939>.
20. Santos ROM, Romano VF, Engstrom EM. Vínculo longitudinal na Saúde da Família: construção fundamentada no modelo de atenção, práticas interpessoais e organização dos serviços. *Physis.* 2018;28(2):e280206. DOI: <https://doi.org/10.1590/S0103-73312018280206>
21. Baratieri T, Lentsck MH, Falavina LP, Soares LG, Prezotto KH, Pitilin EB. Longitudinalidade do cuidado: fatores associados à adesão à consulta puerperal segundo dados do PMAQ-AB. *Cad Saude Publica.* 2022;38(3):e00103221. DOI: <https://doi.org/10.1590/0102-311X00103221>
22. Baratieri T, Chaves ACP, Oliveira IB, Pelazza BB, Lentsck MH, Sangaleti CT, Pitilin EB, Malaquias TSM. Fatores de acesso associados à adesão à consulta pós-parto na atenção primária à saúde. *Revista Contexto & Saúde.* 2024;(48):e14651. DOI: <https://doi.org/10.21527/2176-7114.2024.48.14651>
23. Cruz MM, Souza RBC, Torres RMC, Abreu DMF, Reis AC, Gonçalves AL. Usos do planejamento e autoavaliação nos processos de trabalho das equipes de Saúde da Família na Atenção Básica. *Saude debate.* 2014;38(esp):124-139. DOI: <https://doi.org/10.5935/0103-1104.2014S010>
24. Oliveira JS, Filho JBC. Avaliação da atenção pré-natal na rede básica de saúde em Sergipe – Programa Nacional de Melhoria do Acesso da Qualidade da Atenção Básica PMAQ. *Rev. Rede cuid. Saúde.* 2021;15(1):13-27.
25. Campos GWS. Equipes de referência e apoio especializado matricial: um ensaio sobre reorganização do trabalho em saúde. *Cienc Saude Coletiva.* 1999;4(2):393-403. DOI: <https://doi.org/10.1590/S1413-81231999000200013>
26. Vendruscolo C, Metelski FK, Maffisoni AL, Tesser CD, Trindade LL. Characteristics and performance of professionals of the Expanded Family Health and Basic Healthcare Centers. *Rev Esc Enferm USP.* 2020;54:e03554. DOI: <https://doi.org/10.1590/S1980-220X2018033003554>
27. Melo EA, Almeida PF de, Lima de LD, Giovanella L. Reflections on changes in the federal funding model of Primary Health Care in Brazil. *Saude debate.* 2019;43(5):137-144. DOI: <https://doi.org/10.1590/0103-11042019S512>
28. Marrafon CB, Alcantra DS, Magalhães CCRGN, Buges NM, Melo MP, Costa GD. Referência e contrarreferência na atenção primária. *Cuadernos De Educación Y Desarrollo.* 2024;6(8):e5020. DOI: <https://doi.org/10.55905/cuadv16n8-001>
29. Marini JM, Ronchetti R, Romero SS. Redes de atenção à saúde: a importância da referência e contrarreferência para a atuação integral na medicina. *Revista Perspectiva.* 2024;48(7):49–62. DOI: <https://doi.org/10.31512/persp.v.48.n.7.2024.402.p.49-62>

EVALUATION OF THE WORK PROCESS OF PRIMARY HEALTH CARE TEAMS IN PRENATAL CARE

30. Barbieri MRB, Barildi NG, Fumincelli L, Souza BF, Wernet M, Fabbro MRC. Cuidado pré-natal e integralidade: revisão de escopo. Res Soc Dev. 2021;10(12):e429101220639. DOI: <https://doi.org/10.33448/rsd-v10i12.20639>

Submitted on: May 26, 2023

Accepted on: September 6, 2024

Published on: April 10, 2025

Authors' contributions:

Érica de Brito Pitilin; Conceptualization; Data curation; Formal analysis; Funding acquisition; Investigation; Methodology; Project administration; Resources; Supervision; Validation; Visualization; Writing – original draft, Writing – review & editing.

Daniela Savi Geremia; Conceptualization; Data curation; Formal analysis; Funding acquisition; Investigation; Methodology; Resources; Validation; Visualization; Writing – original draft, Writing – review & editing.

Jéssica Ferreira; Conceptualization; Data curation; Formal analysis; Funding acquisition; Investigation; Methodology; Resources; Validation; Visualization; Writing – original draft, Writing – review & editing.

Yaná Tamara Tomasi; Conceptualization; Data curation; Formal analysis; Funding acquisition; Investigation; Methodology; Resources; Validation; Visualization; Writing – original draft, Writing – review & editing.

Carine Vendruscolo; Conceptualization; Data curation; Formal analysis; Funding acquisition; Investigation; Methodology; Resources; Validation; Visualization; Writing – original draft, Writing – review & editing.

Tatiane Baratieri; Conceptualization; Data curation; Formal analysis; Funding acquisition; Investigation; Methodology; Resources; Validation; Visualization; Writing – original draft, Writing – review & editing.

Maicon Henrique Lentsck; Conceptualization; Data curation; Formal analysis; Funding acquisition; Investigation; Methodology; Resources; Validation; Visualization; Writing – original draft, Writing – review & editing.

Janine Schirmer; Conceptualization; Data curation; Formal analysis; Funding acquisition; Investigation; Methodology; Resources; Validation; Visualization; Writing – original draft, Writing – review & editing.

All authors approved the final version of the text.

Conflict of interest: There is no conflict of interest.

**EVALUATION OF THE WORK PROCESS OF
PRIMARY HEALTH CARE TEAMS IN PRENATAL CARE**

Funded by: Federal University of Fronteira Sul (UFFS).

Corresponding author:

Jéssica Ferreira

Federal University of Fronteira Sul (UFFS).

SC-459, km 2, s/n. Exit to Guatambu – Fronteira Sul – Room 306 – ZIP Code 89812270

Chapecó/SC – Brazil

jessicaferreira2603@gmail.com

Editor-in-chief: Dr. Adriane Cristina Bernat Kolankiewicz

This is an open access article distributed under the terms of the Creative Commons license.

